arXiv:math/9712221v1 [math.GT] 3 Dec 1997

Pure braids, a new subgroup of the mapping class group and finite-type invariants

Jerome Levine
Brandeis University
Waltham, MA 02254 USA
levine@binah.cc.brandeis.edu

February 1, 2008

Abstract

In the study of the relation between the mapping class group Γ_g of a surface Σ_g of genus g and the theory of finite-type invariants of homology 3-spheres, three subgroups of Γ_g play a large role. They are the Torelli group $\mathcal{T}_g$, the Johnson subgroup $\mathcal{K}_g$ and a new subgroup $\mathcal{L}_g^L$, which contains $\mathcal{K}_g$, defined by a choice of a Lagrangian subgroup $L \subseteq H_1(\Sigma_g)$. In this work we determine the quotient $\mathcal{L}_g^L/\mathcal{K}_g \subseteq \Gamma_g/\mathcal{K}_g$, in terms of the precise description of $\Gamma_g/\mathcal{K}_g$ given by Johnson and Morita. We also study the lower central series of $\mathcal{L}_g^L$ and $\mathcal{K}_g$, using some natural imbeddings of the pure braid group in $\mathcal{L}_g^L$ and the theory of finite-type invariants.

Contents

1 Introduction 1

2 Statement of Results 1
- 2.1 The group $\mathcal{L}_g^L$ 1
- 2.2 Imbedding the pure braid group in the mapping class group 4
- 2.3 Applications of the theory of finite-type invariants 7

3 Proof of Theorem 1 9
- 3.1 Construction of $\mathcal{J}$ 9
- 3.2 Proof of (a) 11
- 3.3 Proof of (b) 11
- 3.4 Proof of Corollary 1 12
- 3.5 Proof of Corollary 2 13

---

*Partially supported by NSF grant DMS-96-26639

4 Proof of Theorem 2 **13**
4.1 The case $m = 2$ 14
4.2 The cases $m \geq 3$ 14

5 Proof of Theorems 3 and 4 **15**

6 Proof of Theorems 5, 6 and 7 **15**
6.1 Proof of Theorems 5 and 6 15
6.2 Proof of Theorem 7 17

7 Proof of Theorem 8 and Theorem 9 **18**
7.1 Proof of Theorem 8 18
7.2 Proof of Theorem 9 19

1 Introduction

Let Γ_g denote the group of isotopy classes of orientation-preserving diffeomorphisms of the *bounded* surface Σ_g of genus g with one boundary component which are the identity on the boundary. In [7] and [8] the relation between the mapping class group and the theory of finite-type invariants of homology spheres is investigated. It turns out that the three different notions of finite-type which are discussed in [7] require one to consider three different subgroups of the mapping class group. Two of these subgroups are familiar from previous work of D. Johnson, S. Morita and others on the structure of the mapping class group (see [15]) and [22] for surveys). These are the Torelli group $\mathcal{T}_g$ (maps which induce the identity automorphism on $\mathrm{H}_1(\Sigma_g)$) and the *Johnson subgroup* $\mathcal{K}_g$ (generated by Dehn twists on bounding closed curves). But the original notion of finite-type introduced by Ohtsuki in [25] requires one to consider a new subgroup defined as follows. Choose a *Lagrangian* subspace $L \subseteq H = \mathrm{H}_1(\Sigma_g)$, i.e. a summand of rank g on which the intersection form vanishes. Define $\mathcal{L}_g^L$ to be the subgroup of Γ_g generated by Dehn twists on closed curves whose homology class lies in L. Of course $\mathcal{L}_g^L$ depends on L but its conjugacy class is independent of the choice of L. It is also clear that $\mathcal{K}_g \subseteq \mathcal{L}_g^L$. In fact, the relation between $\mathcal{L}_g^L$ and $\overline{\mathcal{L}_g^L}$ is comparable to the relation between $\mathcal{K}_g$ and $\mathcal{T}_g$.

In this work we will give a precise determination of how $\mathcal{L}_g^L$ sits in Γ_g, or, more precisely, how $\mathcal{L}_g^L/\mathcal{K}_g$ sits in $\Gamma_g/\mathcal{K}_g$, using the description given by Johnson [12] and [13]. The analogous result for a *closed* surface is obtained as a corollary and is, in fact, somewhat simpler to state than the bounded case.

A second purpose of this paper is to study the lower central series of the groups $\mathcal{L}_g^L$ and $\mathcal{K}_g$. We show that the lower central series of $\mathcal{L}_g^L$ is dominated by a refinement of the relative weight filtration of Γ_g. We also make use of a natural imbedding of the pure braid group P_g on g strands into $\mathcal{L}_g^L$ (first used by Oda, in an unpublished paper [24][1] to give lower bounds for the ranks of the associated gradeds of the relative weight filtration of Γ_g), and of P_{2g} into $\mathcal{K}_g$, to give lower bounds on the ranks of the lower central series quotients of these groups. These estimates augment the lower bounds given in [8] using the theory of finite-type invariants of homology 3-spheres.

Of course the definitive results on the lower central series of $\mathcal{T}_g$ are those of Hain [10].

2 Statement of Results

2.1 The group $\mathcal{L}_g^L$

Let $\overline{\mathcal{L}_g^L}$ denote the subgroup of Γ_g consisting of maps whose induced automorphism of H is the identity on L. Then $\mathcal{L}_g^L \subseteq \overline{\mathcal{L}_g^L}$ and $\mathcal{T}_g \subseteq \overline{\mathcal{L}_g^L}$. Under the obvious isomorphism $\Gamma_g/\mathcal{T}_g \cong Sp(\mathbb{Z}, 2g)$, the symplectic group over $\mathbb{Z}$ of genus g, $\overline{\mathcal{L}_g^L}/\mathcal{T}_g$ corresponds to the subgroup B_g consisting of symplectic automorphisms of H which are the identity on L. B_g is isomorphic to $\mathcal{S}_2(L)$, the additive subgroup of symmetric elements of $L \otimes L$. If $f \in B_g$, then $\phi - 1$ induces an element of $\mathrm{Hom}(H/L, L) \cong \mathrm{Hom}(L^*, L) \cong L \otimes L$ (using duality via the intersection pairing on H). Alternatively if we choose a symplectic basis $\{x_i, y_i\}$ for H, where $\{x_i\}$ is a basis of L, then ϕ is represented by a matrix in the form $\left(\begin{smallmatrix} I & A \\ 0 & I \end{smallmatrix}\right)$ and $\phi \longrightarrow A$ defines an isomorphism of B_g with the additive group of symmetric integral matrices. See [7] for more details.

---

[1] The author thanks S. Morita for informing him of the existence of this paper and providing a copy.

Recall the Johnson homomorphism $J : \mathcal{T}_g \longrightarrow \Lambda^3 H$, which is onto and whose kernel is the Johnson subgroup $\mathcal{K}_g$ (see [12] and [13]). If $f \in \mathcal{T}_g$ and $\alpha \in F = \pi_1(\Sigma_g)$, the free group on $2g$ generators, then we can write

$$f_*(\alpha) = \alpha \Phi_f(\alpha) \bmod F_3$$

where, for any group G, G_q denotes the q-th lower central series subgroup: $G_1 = G$, $G_{q+1} = [G, G_q]$. $\Phi_f(\alpha)$ can be regarded, therefore, as an element in $F_2/F_3 \cong \Lambda^2 H$ and so Φ_f defines a homomorphism $H \longrightarrow \Lambda^2 H$ or, dually, an element of $H \otimes \Lambda^2 H$. Johnson shows in [12] that Φ_f lies in the subgroup $\Lambda^3 H \subseteq H \otimes \Lambda^2 H$, where this imbedding is defined by

$$h_1 \wedge h_2 \wedge h_3 \longrightarrow h_1 \otimes (h_2 \wedge h_3) + h_2 \otimes (h_3 \wedge h_1) + h_3 \otimes (h_1 \wedge h_2)$$

and, in [13], that the correspondence $f \longrightarrow \Phi_f$ defines an *isomorphism* $\mathcal{T}_g/\mathcal{K}_g \cong \Lambda^3 H$. From this we can, therefore, regard $\overline{\mathcal{L}_g^L}/\mathcal{K}_g$ as an extension of $\Lambda^3 H$ by B_g.

Theorem 1. *There exists a homomorphism*

$$\mathcal{J} : \overline{\mathcal{L}_g^L} \longrightarrow \Lambda^3(H/L) \oplus H/L \tag{1}$$

extending the Johnson homomorphism, which satisfies the following:

- *(a)* $\mathcal{J}$ *is onto,*
- *(b)* $\ker \mathcal{J} = \mathcal{L}_g^L$
- *(c) There is a commutative diagram*

$$\begin{tikzcd}
 & & B_g & 1 & \\
1 \arrow[r] & \mathcal{L}_g^L \arrow[r] & \overline{\mathcal{L}_g^L} \arrow[u] \arrow[r, "\mathcal{J}"] & \Lambda^3(H/L) \oplus H/L \arrow[u] \arrow[r] & 1 \\
1 \arrow[r] & \mathcal{K}_g \arrow[u] \arrow[r] & \mathcal{T}_g \arrow[u] \arrow[r, "J"'] & \Lambda^3 H \arrow[u, "p \oplus c"] \arrow[r] & 1 \\
 & 1 \arrow[u] & 1 \arrow[u] & &
\end{tikzcd} \tag{2}$$

where p is the projection $\Lambda^3 H \longrightarrow \Lambda^3(H/L)$ and c is the composition of the projection $H \longrightarrow H/L$ with the contraction map $\Lambda^3 H \longrightarrow H$ defined by

$$h_1 \wedge h_2 \wedge h_3 \longrightarrow (h_1 \cdot h_2)h_3 + (h_2 \cdot h_3)h_1 + (h_3 \cdot h_1)h_2$$

If $\Gamma_{g,0}$ denotes the mapping class group of a *closed* surface $\Sigma_{g,0}$ of genus g, i.e. the group of isotopy clases of orientation-preserving diffeomorphisms of $\Sigma_{g,0}$, then we can define the analogous subgroups $\overline{\mathcal{L}^L_{g,0}}, \mathcal{L}^L_{g,0} \subseteq \Gamma_{g,0}$. We also have the corresponding Torelli group $\mathcal{T}_{g,0}$ and Johnson subgroup $\mathcal{K}_{g,0}$. The homomorphism $J : \mathcal{T}_g \longrightarrow \Lambda^3 H$ is shown, in [12], to induce a homomophism $J' : \mathcal{T}_{g,0} \longrightarrow \Lambda^3 H/H$ whose kernel is $\mathcal{K}_{g,0}$.

Corollary 1. *There exists an epimorphism* $J_0 : \overline{\mathcal{L}^L_{g,0}} \longrightarrow \Lambda^3(H/L)$ *whose kernel is* $\mathcal{L}^L_{g,0}$ *and which satisfies the commutative diagram:*

$$\begin{tikzcd}
1 \arrow[r] & \mathcal{L}^L_{g,0} \arrow[r] & \overline{\mathcal{L}^L_{g,0}} \arrow[r, "J_0"] & \Lambda^3(H/L) \arrow[r] & 1 \\
1 \arrow[r] & \mathcal{K}_{g,0} \arrow[r] \arrow[u] & \mathcal{T}_{g,0} \arrow[r, "J'"] \arrow[u] & \Lambda^3 H/H \arrow[r] \arrow[u, "p'"] & 1
\end{tikzcd} \tag{3}$$

where p' is induced by p.

We can also characterize $\mathcal{L}^L_{g,0} \subseteq \Gamma_{g,0}$ in terms of the induced outer automorphisms of $\pi = \pi_1(\Sigma_{g,0})$. let $\pi_L \subseteq \pi$ be the subgroup of elements whose homology class belongs to L. If $f \in \overline{\mathcal{L}^L_{g,0}}$, then $f_*|\pi_L \equiv \mathrm{id} \mod \pi_2$.

Corollary 2. *If* $f \in \Gamma_{g,0}$*, then* $f \in \mathcal{L}^L_{g,0}$ *if and only if* $f_*|\pi_L \equiv id \mod [\pi, \pi_L]$.

Note that this condition on f_* depends only on its outer automorphism class.

We also identify the lower central series terms of $\mathcal{L}^L_g$, modulo $\mathcal{K}_g$. Let $K = \ker(p \oplus e)$. By Theorem 1, we have an isomorphism

$$\mathcal{J}' : \mathcal{L}^L_g \cap \mathcal{T}_g / \mathcal{K}_g \cong K \tag{4}$$

We now define a filtration of $\Lambda^3 H$:

$$\Lambda^3 H = K_0 \supseteq K_1 \supseteq K_2 \supseteq K_3 = \Lambda^3 L \supseteq K_4 = 0$$

by letting K_m be the subgroup of $\Lambda^3 H$ generated by elements $h_1 \wedge h_2 \wedge h_3$, where at least m of the h_i lie in L. Note that $K_1 \supsetneq K \supsetneq K_2$.

Theorem 2. *Under the isomorphism $\mathcal{J}'$ of* (4) *the lower central series terms of $\mathcal{L}^L_g$ satisfy:*

$$((\mathcal{L}^L_g)_m \cdot \mathcal{K}_g)/\mathcal{K}_g \cong K_m \textit{ for } m \geq 2$$

In particular $(\mathcal{L}^L_g)_4 \subseteq \mathcal{K}_g$*, but* $(\mathcal{L}^L_g)_3 \nsubseteq \mathcal{K}_g$.

Furthermore $\mathcal{L}^L_g \cap \mathcal{T}_g = \mathcal{K}_g \cdot [\mathcal{L}^L_g, \mathcal{T}_g]$.

Question 1. Does $[\mathcal{L}^L_g, \mathcal{L}^L_g]$ contain $\mathcal{K}_g$?

We obtain some further information on the lower central series of $\mathcal{L}_g^L$ inside $\mathcal{K}_g$. Recall the filtration on Γ_g defined by:

$$\Gamma_g[n] = \{f \in \Gamma_g | f_* : F \longrightarrow F \text{ is the identity mod } F_{n+1}\}$$

This is called the *relative weight filtration* in [24]. Thus $\Gamma_g[1] = \mathcal{T}_g$ and $\Gamma_g[2] = \mathcal{K}_g$.

Theorem 3. *$(\mathcal{L}_g^L)_q \subseteq \Gamma_g[n]$ if $q \geq \binom{n+3}{2} - 6$*

Thus $\mathcal{L}_g^L$ is residually nilpotent.

In fact we have a more delicate description of the relation between the lower central series of $\mathcal{L}_g^L$ and the relative weight filtration which requires us to define a more refined filtration. We first recall the extended Johnson homomorphisms from [15] as formulated by Morita [20]. Let $\mathbb{L}(H)$ be the graded free Lie algebra (over $\mathbb{Z}$) on H. $\mathbb{L}_n(H)$ is generated by the brackets of length n. Then

$$J_n : \Gamma_g[n] \longrightarrow H \otimes \mathbb{L}_{n+1}(H)$$

is defined by $f_*(\alpha) = \alpha J_n(\alpha) \bmod F_{n+2}$, where $J_n(\alpha) \in F_{n+1}/F_{n+2} \cong \mathbb{L}_{n+1}(H)$. Then $\Gamma_g[n+1] = \ker J_n$ and

$$\operatorname{Im} J_n \subseteq \ker\{b : H \otimes \mathbb{L}_{n+1}(H) \longrightarrow \mathbb{L}_{n+2}(H)\} \tag{5}$$

where $b(h \otimes \alpha) = [h, \alpha]$ (see [21]). Thus $J_1((\mathcal{L}_g^L)_m) = K_m$ (by Theorem 2).

We generalize the filtration $\{K_m\}$ as follows:

$$\mathbb{L}_m(H) = \mathbb{L}_m^0(H) \supseteq \mathbb{L}_m^1(H) \supseteq \cdots \supseteq \mathbb{L}_m^m(H) = \mathbb{L}_m(L) \supseteq \mathbb{L}_m^{m+1}(H) = 0$$

where $\mathbb{L}_m^r(H)$ is the subgroup generated by brackets of length m of which at least r of the entries belong to L. We can then define a filtration of $H \otimes \mathbb{L}_m(H)$ by $\mathcal{F}_m^r = (L \otimes \mathbb{L}_m^{r-1}(H)) \oplus (H \otimes \mathbb{L}_m^r(H))$. Thus $\mathcal{F}_m^{m+1} = L \otimes \mathbb{L}_m(L)$ and $\mathcal{F}_m^{m+2} = 0$.

Theorem 4. *$J_n((\mathcal{L}_g^L)_{q+r}) \subseteq \mathcal{F}_{n+1}^r$ if $q \geq \binom{n+3}{2} - 6$.*

2.2 Imbedding the pure braid group in the mapping class group

It has been pointed out several times in the literature that one can define interesting maps from the group of *framed* pure braids to the mapping class group. The framed pure braid group on h strands, $\mathcal{P}_h$, which is canonically isomorphic to the Cartesian product of the usual pure braid group $\mathcal{P}_h^\circ$ with the free abelian group of rank h, can be represented as the group of isotopy classes of diffeomorphisms of the disk with h holes, D_h, which are the identity on the boundary. Any imbedding of D_h into Σ_g defines, therefore, a homomorphism $\theta : \mathcal{P}_h \longrightarrow \Gamma_g$. For example one can imbed D_{2g} into Σ_g as in Figure 1. This map was considered by Hatcher-Thurston [11] and, with a formulation in terms of tangles, Matveev-Polyak [18] in a slightly more general setting of what Matveev-Polyak call *admissible* braids. It is not hard to see that $\theta(\mathcal{P}_{2g}) \subseteq \mathcal{L}_g^L$, where L is the subspace spanned by the meridians of the handles. We



Figure 1. The Hatcher-Thurston map

will be more interested, though, in a smaller version of this map. Consider the imbedding $D_g \subseteq \Sigma_g$ indicated in Figure 2. The resulting map $\psi : \mathcal{P}_g \longrightarrow \Gamma_g$ was considered in [24] Oda who showed that ψ induces *imbeddings* of the lower-central series quotients of $\mathcal{P}_g$ into the associated graded quotients of Γ_g using the relative weight filtration: $\psi_n : (\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \longrightarrow \Gamma_g[n-1]/\Gamma_g[n]$. The well-known split exact sequence $1 \longrightarrow F^{g-1} \longrightarrow \mathcal{P}_g \longrightarrow \mathcal{P}_{g-1} \longrightarrow 1$ gives an isomorphism:

$$(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \cong (\mathcal{P}_{g-1})_n/(\mathcal{P}_{g-1})_{n+1} \oplus F^{g-1}_n/F^{g-1}_{n+1} \tag{6}$$

See e.g. [2].This enables Oda to use ψ_n to give some explicit lower bounds for the rank of $\Gamma_g[n-1]/\Gamma_g[n]$.

We observe easily that ψ actually maps into $\mathcal{L}_g^L$, for the correct choice of L, and describe how the image fits into the refined filtration $\mathcal{F}^r_m$ of $H \otimes \mathbb{L}(H)$.

Theorem 5. *$\psi(\mathcal{P}_g) \subseteq \mathcal{L}_g^L$ and $\mathrm{Im}(J_{n-1}\psi_n) \subseteq \mathcal{F}^{n+1}_n = L \otimes \mathbb{L}_n(L)$.*

Thus $(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \subseteq \Gamma_g[n-1]/\Gamma_g[n]$ lies in the *bottom stage* of this filtration.

Remark 2.1. It is not hard to prove, by a similar argument, that the conclusions of Theorem 5 are also true for the Hatcher-Thurston map.

As in Equation (5) $\mathrm{Im}(J_{n-1}\psi_n) \subseteq \ker\{b : L \otimes \mathbb{L}_n(L) \longrightarrow \mathbb{L}_{n+1}(L)\}$, but for $n \geq 3$, this inclusion is definitely proper. One sees this by computing the difference between the rank of $\ker b$ and the rank of $(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1}$, using Equation (6). For example, for $n = 3$ this difference is $\frac{1}{6}(g^3 - g)$ and for $n = 4$ it is $\frac{1}{8}(g^3 - g)(g - 2)$. For $n \leq 2$ see Theorem 6 below.

The non-triviality of $J_{n-1}\psi_n$ shows that $\mathcal{L}_g^L$ is not nilpotent. More precisely, combining this with Theorem 3 gives a lower bound on the rate of descent of the lower central series of $\mathcal{L}_g^L$. We denote the rank of $(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1}$ by $r(g,n)$. This can be computed explicitly from Equation (6).

Corollary 3. *The image of the map $(\mathcal{P}_g)_n/(\mathcal{P}_g)_q \longrightarrow (\mathcal{L}_g^L)_n/(\mathcal{L}_g^L)_q$ has rank $\geq r(g,n)$ if $q \geq \binom{n+3}{2} - 6$.*



Figure 2. Imbedding of $\mathcal{P}_g$ into $\mathcal{L}_g^L$

This estimate is independent of the estimate on the rank of the lower central series of $\mathcal{L}_g^L$ given in [8]. Recall the map $\phi_n^L : (\mathcal{L}_g^L)_{3n}/(\mathcal{L}_g^L)_{3n+1} \otimes \mathbb{Q} \longrightarrow \mathcal{A}_n^{\mathrm{conn}}(\emptyset)$ defined in [8] and shown there (Theorem 7) to be onto if $g \geq 5n+1$, where $\mathcal{A}_n^{\mathrm{conn}}(\emptyset)$ is a vector space defined by trivalent graphs with $2n$ vertices and $3n$ edges. Now it is easy to see that the composition $(\mathcal{P}_g)_{3n} \longrightarrow (\mathcal{L}_g^L)_{3n} \longrightarrow \mathcal{A}_n^{\mathrm{conn}}(\emptyset)$ is zero. If $h \in \operatorname{Im}\psi$ then $S_h^3 = S^3$ since h extends to a diffeomorphism of the handlebody bounded by Σ_g. Thus combining Corollary 3 and [8, Theorem 7] we have:

Corollary 4. $\operatorname{rank}(\mathcal{L}_g^L)_{3n}/(\mathcal{L}_g^L)_q \geq r(g,3n) + \dim \mathcal{A}_n^{conn}(\emptyset)$ *if* $q \geq \binom{3n+3}{2} - 6$ *and* $g \geq 5n+1$.

For the special cases $n = 1, 2$ we have:

Theorem 6. *(a)* ψ_1 *induces an isomorphism* $\mathcal{P}_g/(\mathcal{P}_g)_2 \cong \mathcal{L}_g^L/(\mathcal{L}_g^L \cap \mathcal{T}_g) \cong B_g$.

(b) $J_1\psi_2$ *induces an isomorphism* $(\mathcal{P}_g)_2/(\mathcal{P}_g)_3 \otimes \mathbb{Q} \cong ((\mathcal{L}_g^L)_3 \cdot \mathcal{K}_g)/\mathcal{K}_g \otimes \mathbb{Q} \cong \Lambda^3 L \otimes \mathbb{Q}$. *Thus* $\psi^{-1}(\mathcal{K}_g) = (\mathcal{P}_g)_3$.

Question 2. Can (b) be improved to say $(\mathcal{P}_g)_2/(\mathcal{P}_g)_3 \cong \Lambda^3 L$?

Remark 2.2. For the Hatcher-Thurston map θ one has $J_1\theta((\mathcal{P}_{2g})_2) \subseteq K_3 = \Lambda^3 L$, according to Remark 2.1, but it is also not hard to show that $J_1(\theta(\mathcal{P}_{2g}) \cap \mathcal{T}_g) \subseteq K_2$. A consequence of this is that $\theta(\mathcal{P}_{2g}) \subsetneq \mathcal{L}_g^L$.

We will consider one more mapping of the pure braid group into the mapping class group. Consider the imbedding $D_g \subseteq \Sigma_g$ indicated in Figure 3. Denote the resulting map by $\kappa : \mathcal{P}_g \longrightarrow \Gamma_g$. We will prove:



Figure 3. Imbedding of $\mathcal{P}_g$ into $\mathcal{K}_g$

Theorem 7. *$\kappa(\mathcal{P}_g) \subseteq \mathcal{K}_g$ and $\kappa((\mathcal{P}_g)_n) \subseteq \Gamma_g[2n]$. The induced map $\kappa_n : (\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \longrightarrow \Gamma_g[2n]/\Gamma_g[2n{+}1]$ is a monomorphism. Thus the induced map $(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \longrightarrow (\mathcal{K}_g)_n/(\mathcal{K}_g)_{n+1}$ is also a monomorphism and so $\operatorname{rank}(\mathcal{K}_g)_n/(\mathcal{K}_g)_{n+1} \geq r(g,n)$.*

As above we have, from [8], a map $\phi_n^K : (\mathcal{K}_g)_n/(\mathcal{K}_g)_{n+1} \longrightarrow \mathcal{A}_n^{\text{conn}}(\emptyset)$ which is onto if $g \geq 5n+1$. The composition $(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \longrightarrow (\mathcal{K}_g)_n/(\mathcal{K}_g)_{n+1} \longrightarrow \mathcal{A}_n^{\text{conn}}(\emptyset)$ is zero since, if $h \in \operatorname{Im} \phi_n^K$ then $S_h^3 = S^3$. Thus we have, from Theorem 7 and [8]:

Corollary 5. *$\operatorname{rank}(\mathcal{K}_g)_n/(\mathcal{K}_g)_{n+1} \geq r(g,n) + \dim \mathcal{A}_n^{conn}(\emptyset)$ if $g \geq 5n+1$.*

2.3 Applications of the theory of finite-type invariants

The fact that there are useful connections between the theory of finite-type invariants of homology spheres and the structure of subgroups of the mapping class group was established in [7] and [8]. In particular, as we have mentioned above, Theorem 6 of [8] gives lower bounds for the dimensions of the graded quotients of the lower central series of the subgroups $\mathcal{L}_g^L, \mathcal{T}_g$ and $\mathcal{K}_g$ in terms of certain vector spaces $\mathcal{A}_n^{\text{conn}}(\emptyset)$ defined by connected trivalent graphs.

As a further application of this result we have the following relation between the lower central series of these groups.

Theorem 8. *Let $g \geq 5n+1$. Then*

1. $(\mathcal{K}_g)_n \nsubseteq (\mathcal{T}_g)_{2n+1} \cup (\mathcal{L}_g^L)_{3n+1}$,
2. $(\mathcal{T}_g)_{2n} \nsubseteq (\mathcal{K}_g)_{n+1} \cup (\mathcal{L}_g^L)_{3n+1}$,
3. $(\mathcal{L}_g^L)_{3n} \nsubseteq (\mathcal{T}_g)_{2n+1} \cup (\mathcal{K}_g)_{n+1}$.

We also examine the map $\psi : \mathcal{P}_g \longrightarrow \mathcal{L}_g^L$, defined in the previous section, and its relationship to the theory of finite-type invariants of knots and homology spheres. Let $\mathcal{V}$ denote the $\mathbb{Q}$-vector space generated by isotopy classes of oriented smooth knots in S^3 and let $\mathcal{M}$ denote the $\mathbb{Q}$-vector space generated by diffeomorphism classes of smooth oriented homology spheres. In [3] Garoufalidis defines a natural linear map $\Psi : \mathcal{V} \longrightarrow \mathcal{M}$ by $\Psi(K) = S^3_K$, where, for any knot K in a homology sphere M, M_K denotes the homology sphere obtained by doing a $+1$-surgery on $K \subseteq M$. Filtrations are defined on $\mathcal{V}$ and $\mathcal{M}$ by the following devices. For any positive integer q, a *singular q-knot* is an immersion of S^1 into S^3 with exactly q transverse (ordered) double points. If K is a singular q-knot, the resolution $\hat{K}$ of K is the element of $\mathcal{V}$ defined by $\hat{K} = \sum_\epsilon (-1)^{|\epsilon|} K_\epsilon$, where ϵ ranges over all sequences $(\epsilon_1, \ldots, \epsilon_q), \epsilon_i = 0$ or 1, $|\epsilon| = \sum_i \epsilon_i$ and K_ϵ denotes the knot obtained from K by resolving each double point to give a positive, resp. negative crossing according to whether ϵ_i is, resp. 0 or 1. Now define $\mathcal{F}_q(\mathcal{V})$ to be the subspace of $\mathcal{V}$ generated by the resolutions of all q-singular knots. This gives a decreasing filtration of $\mathcal{V}$ (see [1]). To deal with $\mathcal{M}$ define, for any q-component link J in a homology sphere, an element $[M; J] \in \mathcal{M}$ by the formula $[M; J] = \sum_\epsilon (-1)^{|\epsilon|} M_{J_\epsilon}$ where ϵ is as above and J_ϵ is the sublink of J consisting of exactly those components for which $\epsilon_i = 1$. Now $\mathcal{F}_q(\mathcal{M})$ is defined to be the subspace of $\mathcal{M}$ generated by all $[M; J]$ for which J has q components. This is a decreasing filtration of $\mathcal{M}$ (see [25]). It is shown in [25] and [5] that $\mathcal{F}_{3q-2}(\mathcal{M}) = \mathcal{F}_{3q}(\mathcal{M})$, for all q.

The associated graded quotients

$$\mathcal{G}_q(\mathcal{V}) = \mathcal{F}_q(\mathcal{V})/\mathcal{F}_{q+1}(\mathcal{V}) \text{ and } \mathcal{G}_q(\mathcal{M}) = \mathcal{F}_{3q}(\mathcal{M})/\mathcal{F}_{3q+1}(\mathcal{M})$$

are described in terms of spaces of trivalent graphs $\mathcal{A}(S^1), \mathcal{A}(\emptyset)$, where $\mathcal{A}(X)$, for any 1-manifold X, is a certain $\mathbb{Q}$-vector space defined from trivalent graphs which contain X as a prescribed subgraph (see [1], [25] and [16]). It was conjectured by Garoufalidis in [3] and proved by Habegger [9] that $\Psi(\mathcal{F}_{2q}(\mathcal{V})) \subseteq \mathcal{F}_{3q}(\mathcal{M})$ and so we have induced maps $\Psi_q : \mathcal{G}_{2q}(\mathcal{V}) \longrightarrow \mathcal{G}_q(\mathcal{M})$. This map is studied in [4], using the work of [16], and shown to be non-trivial for all q.

The map $\psi : \mathcal{P}_g \longrightarrow \mathcal{L}^L_g$ and $\Psi : \mathcal{V} \longrightarrow \mathcal{M}$ are related by means of 'actions' of $\mathcal{P}^\circ_g$ on $\mathcal{V}$ and $\mathcal{L}^L_g$ on $\mathcal{M}$. Let K be an oriented knot in S^3 and $D \subseteq S^3$ an imbedded disk which meets K transversely in g points so that, at each intersection point, K points in the same normal direction to D. If b is any g-strand pure braid then we can cut S^3 along D and insert b so that the orientations agree, to form a new knot K_b. This action was defined by Stanford in [26] who showed that if $b \in (\mathcal{P}^\circ_g)_m$, then $K_b - K \in \mathcal{F}_m(\mathcal{V})$. Note also that if b is a framed pure braid and K is a framed knot, then K_b is also framed in a natural way. If s is the self-linking number of K and $\{s_i\}$ are the self-linking numbers of the strands of b, then the self-linking number of K_b is $s + \sum_i s_i$.

We now describe the action of $\mathcal{L}^L_g$ on $\mathcal{M}$. Suppose M is a homology sphere and $i : \Sigma_g \subseteq M$ is an imbedding. If $L_i = L \subseteq H = \mathrm{H}_1(\Sigma_g)$ is the kernel of the inclusion into one of the complementary summands, then, for any $f \in \mathcal{L}^L_g$, we can cut M open along $i(\Sigma_g)$ and reglue, using f, to define a new homology sphere M_f (see [7]). It is shown in [7] that if $f \in (\mathcal{L}^L_g)_m$ then $M_f - M \in \mathcal{F}_m(\mathcal{M})$.

Now suppose we are given K, D as above and $b \in \mathcal{P}^\circ_g$. Choose an imbedding $i : \Sigma_g \longrightarrow S^3$ whose image is the boundary of a regular neighborhood of $K \cup D$. Then i will also define an imbedding j of Σ_g into S^3_K. Let b denote also the framed braid whose self-linking numbers are determined by the equations $\sum_j l_{ij} = 0$, where l_{ij} denotes the linking number of the i-th and j-th strand, if $i \neq j$, and the self-linking number of the i-th strand if $i = j$. Then for every $i = 1, \cdots g$, then we have:

Theorem 9. *$\Psi(K_b) = M_{\psi(b)}$, where $M = S^3_K$.*

3 Proof of Theorem 1

As mentioned already, different choices for L result in conjugate subgroups $\mathcal{L}^L_g$ (and also $\overline{\mathcal{L}^L_g}$). Therefore it will suffice to prove Theorems 1-4 for any convenient choice of L. (The other theorems will require certain choices for L). In particular we will choose L to be the subgroup of H generated by meridians of the handles for some representation of Σ_g as the boundary of a handle-body. Thus we can choose a basis $\{x_i, y_i\}$ for $F = \pi_1(\Sigma_g)$ to satisfy:

- The induced basis of H (which we will also denote by x_i, y_i) is *symplectic* ,i.e. under the intersection pairing on H, we have $x_i \cdot x_j = y_i \cdot y_j = 0$ and $x_i \cdot y_j = \delta_{ij}$,
- $\{x_i\}$ is a basis for L,
- The element $[x_1, y_1] \cdots [x_g, y_g] \in F$ represents the boundary curve of Σ_g.

We will call such a basis *admissible for* L, We will also use this term for the induced basis of H.

3.1 Construction of $\mathcal{J}$

We begin by defining a crossed homomorphism on $\overline{\mathcal{L}^L_g}$. Let $F_L \subseteq F$ be the set of all elements of F which map into $L \subseteq H$ under abelianization. Let $\hat{L}$ be the abelianization of F_L. We define

$$\hat{J} : \overline{\mathcal{L}^L_g} \longrightarrow \mathrm{Hom}(\hat{L}, \Lambda^2 H)$$

by the formula $\hat{J}(f)\beta \equiv \alpha^{-1} f_*(\alpha) \mod F_3$, under the identification $\mathcal{F}_2/F_3 \cong \Lambda^2 H$, where $\alpha \in F_L$ is any lift of $\beta \in \hat{L}$.

We then have the following crossed-homomorphism formula

$$\hat{J}(fg) = \hat{J}(g) + \hat{J}(f) \circ g_* \tag{7}$$

where g_* is the automorphism of $\hat{L}$ defined by g.

Composing $\hat{J}$ with the projection $\Lambda^2 H \longrightarrow \Lambda^2(H/L)$ defines

$$\bar{J} : \overline{\mathcal{L}^L_g} \longrightarrow \mathrm{Hom}(\hat{L}, \Lambda^2(H/L))$$

Composing $\hat{J}$ with the contraction map $c : \Lambda^2 H \longrightarrow \mathbb{Z}$ defines

$$J_\omega : \overline{\mathcal{L}^L_g} \longrightarrow \mathrm{Hom}(\hat{L}, \mathbb{Z})$$

c is defined, using the intersection pairing on H, by $h_1 \wedge h_2 \longrightarrow h_1 \cdot h_2$.

We point out two properties of $\bar{J}$ and J_ω.

Lemma 3.1. *(a) $\bar{J}(f)|F_2 = J_\omega(f)|F_2 = 0$, for any $f \in \overline{\mathcal{L}_g^L}$,*

(b) $\bar{J}$ and J_ω are homomorphisms.

Thus we have homomorphisms:

$$\bar{J} : \overline{\mathcal{L}_g^L} \longrightarrow \mathrm{Hom}(L, \Lambda^2(H/L)), \quad J_\omega : \overline{\mathcal{L}_g^L} \longrightarrow \mathrm{Hom}(L, \mathbb{Z})$$

Proof of Lemma 3.1. To prove (a) consider a generating element $[h_1, h_2] \in F_2$. If $f \in \overline{\mathcal{L}_g^L}$, then we can write $f_*(h_i) = h_i l_i$, for some elements $l_i \in F_L$ and so we have:

$$f_*([h_1, h_2]) = [h_1, h_2][l_1, h_2][h_1, l_2][l_1, l_2] \mod F_3$$

Thus $\hat{J}(f)[h_1, h_2] = l_1 \wedge h_2 + h_1 \wedge l_2 + l_1 \wedge l_2$. Since this element projects to zero in $\Lambda^2(H/L)$, (a) follows for $\bar{J}$. For J_ω we note that $l_1 \cdot l_2 = 0$, since the intersection form is zero on L and $h_1 \cdot l_2 + l_1 \cdot h_2 = 0$ since f_* is an isometry.

To prove (b) we need only note that g_* is the identity on L and apply (a) and Equation (7). □

Now we follow Johnson and reformulate $\bar{J}$ first as a homomorphism

$$\tilde{J} : \overline{\mathcal{L}_g^L} \longrightarrow H/L \otimes \Lambda^2(H/L)$$

using the duality isomorphism $\mathrm{Hom}(L, \mathbb{Z}) \cong H/L$ defined by the intersection form on H. If $\{x_i, y_i\}$ is an admissible basis of F for L, then $\tilde{J}(f) = \sum_i y_i \otimes \bar{J}(f) \cdot x_i$.

We have, for any free-abelian group V, a short exact sequence:

$$0 \longrightarrow \Lambda^3 V \xrightarrow{\eta} V \otimes \Lambda^2 V \xrightarrow{\theta} \mathbb{L}_3(V) \longrightarrow 0$$

where $\eta(v_1 \wedge v_2 \wedge v_3) = v_1 \otimes (v_2 \wedge v_3) + v_2 \otimes (v_3 \wedge v_1) + v_3 \otimes (v_1 \wedge v_2)$ and $\theta(v_1 \otimes (v_2 \wedge v_3) = [v_1, [v_2, v_3]]$. We leave the proof as an exercise for the reader.

We can use this sequence to show that $\tilde{J}(\overline{\mathcal{L}_g^L}) \subseteq \Lambda^3(H/L)$. If $f \in \overline{\mathcal{L}_g^L}$, then we can write $f_*(x_i) = x_i C_i$, for some $C_i \in F_2$ which represents $\bar{J}(f) \cdot x_i$, and $f_*(y_i) = y_i l_i$ for some $l_i \in F_L$. By distributivity of brackets we have:

$$f_*[x_i, y_i] \equiv [x_i, y_i][C_i, y_i][x_i, l_i]\alpha_i \mod F_4 \tag{8}$$

where α_i is a product of elements of $\mathcal{F}_3/\mathcal{F}_4 \cong \mathbb{L}_3(H)$ which vanish when projected into $\mathbb{L}_3(H/L)$. Since any element of Γ_g leaves the boundary curve of Σ_g fixed, we have that, up to conjugacy, $f_*(\prod_i [x_i, y_i]) = \prod_i [x_i, y_i]$. Our first deduction from Equation (8) is that $\prod_i [x_i, l_i] \in F_3$. Now if we write out $l_i = (\prod_j x_j^{e_{ij}})C_i'$, for some $C_i' \in F_2$, we can conclude that $e_{ij} = e_{ji}$ and so $\prod_i [x_i, l_i]$ $(\mathrm{mod} F_4)$ reduces to a product of triple brackets, each of which has some x_j as one of its entries, and so vanishes in $\mathbb{L}_3(H/L)$. Thus we can conclude, from Equation (8), that $\sum_i [C_i, y_i] = 0$ in $\mathbb{L}_3(H/L)$. But this means that $\theta \tilde{J}(f) = 0$ and so $\tilde{J}$ actually gives us a homomorphism $\overline{\mathcal{L}_g^L} \longrightarrow \Lambda^3(H/L)$.

We can now set $\mathcal{J} = \tilde{J} \oplus J_\omega$, where J_ω is reformulated as a homomorphism $J_\omega : \overline{\mathcal{L}_g^L} \longrightarrow H/L$, using the duality isomorphism $\mathrm{Hom}(L, \mathbb{Z}) \cong H/L$..

3.2 Proof of (a)

Since the Johnson homomorphism J is onto $\Lambda^3 H$, it is obvious that $\tilde{J}$ is onto. To deal with J_ω we first point out that $J_\omega|\mathcal{T}_g$ can be alternatively defined by the formula: $J_\omega(f) = \sigma J(f)$, where $\sigma : \Lambda^3 H \longrightarrow H \longrightarrow H/L$ is defined by contraction:

$$\sigma(h_1 \wedge h_2 \wedge h_3) = (h_1 \cdot h_2)h_3 + (h_2 \cdot h_3)h_1 + (h_3 \cdot h_1)h_2$$

Let $\{x_i, y_i\}$ be an admissible basis of H. Now choose $f \in \mathcal{T}_g$ so that $J(f) = x_i \wedge y_i \wedge y_j$, for a prescribed $i \neq j$. Then $J_\omega(f) = y_j$ while $\tilde{J}(f) = 0$ in $\Lambda^3(H/L)$. From this it follows that $\mathcal{J}$ is onto. □

3.3 Proof of (b)

It is essentially proved in [7] that $\tilde{J}(\mathcal{L}_g^L) = 0$ but we take a different approach to give a unified proof that $\mathcal{J}(\mathcal{L}_g^L) = 0$.

Suppose that f is a Dehn twist along the curve γ in Σ_g. If γ bounds in Σ_g, then $f \in \mathcal{K}_g$ and there is nothing to prove. If γ does not bound, then we may assume that γ is a meridian curve of a handle in some representation of Σ_g as the boundary of a handle-body. Let $\{x_i, y_i\}$ be the standard basis of F, where x_i represents the meridian of the i-th handle and y_i represents the longitude. Note that this is not necessarily an admissible basis. We may assume that γ represents x_1. Then we see easily that

$$f_*(x_i) = x_i \ (1 \leq i \leq g), \ f_*(y_i) = \begin{cases} y_i & \text{if } 1 \leq i \leq g \\ y_1 x_1 & \text{if } i = 1 \end{cases}$$

All we know about L, though, is that $x_1 \in F_L$ and, as a consequence, an element $w \in F$ lies in F_L only if the exponent sum of y_1 is 0. From this it follows that for any $w \in F_L$, $\tilde{J}(f) \cdot w = a \wedge x_1$, where $a \in H$ is a linear combination of $x_1, \cdots, x_g, y_2, \cdots, y_g$. But then it is clear that $a \wedge x_1$ maps to 0 in $\Lambda^2(H/L)$ and $c(a \wedge x_1) = 0$. Thus $\mathcal{L}_g^L \subseteq \ker \mathcal{J}$

Now suppose $f \in \overline{\mathcal{L}_g^L}$ and $\mathcal{J}(f) = 0$. We want to show that $f \in \mathcal{L}_g^L$. We first recall that the images of $\overline{\mathcal{L}_g^L}$ and of $\mathcal{L}_g^L$ under the canonical map $\Gamma_g \longrightarrow Sp(\mathbb{Z}, g)$ coincide and are equal to the subgroup $B_g \subseteq Sp(\mathbb{Z}, g)$ defined in Section 2.1. This is proved in [7]. Since there exists some $g \in \mathcal{L}_g^L$ so that $f_* = g_*$ on H, then, by replacing f by fg^{-1}, we may assume that $f \in \mathcal{T}_g$. Since $\mathcal{K}_g \subseteq \mathcal{L}_g^L$ it will suffice to find $g \in \mathcal{L}_g^L \cap \mathcal{T}_g$ so that $J(g) = J(f)$. In fact we will find such a $g \in [\mathcal{L}_g^L, \mathcal{T}_g] \subseteq \mathcal{L}_g^L \cap \mathcal{T}_g$ (since $\mathcal{L}_g^L$ is invariant under conjugation by $\mathcal{T}_g$), which will then, in addition, show that $\mathcal{L}_g^L \cap \mathcal{T}_g = [\mathcal{L}_g^L, \mathcal{T}_g] \cdot \mathcal{K}_g$ (part of the conclusion of Theorem 2). Let $K = \ker\{p \oplus c : \Lambda^3 H \longrightarrow \Lambda^3(H/L) \oplus H/L\}$. We need to show that, for any $a \in K$, there exists $g \in [\mathcal{L}_g^L, \mathcal{T}_g]$ such that $J(g) = a$. Some examples of elements of K are, in terms of an admissible basis of H:

1. $a = x_i \wedge x_j \wedge x_k$ for any i, j, k,
2. $a = x_i \wedge x_j \wedge y_k$ for any i, j, k,
3. $a = x_i \wedge y_j \wedge y_k$ for i, j, k distinct and

4. $a = y_i \wedge x_i \wedge y_k + x_j \wedge y_j \wedge y_k$ for i, j, k distinct.

In fact we will leave it as an exercise for the reader to prove that any element of K is a linear combination of elements of these four types.

To compute J on $[\mathcal{L}_g^L, \mathcal{T}_g]$ we need the following observation. Let $f \in \Gamma_g$ and $h \in \mathcal{T}_g$. Then we have, for the classical Johnson homomorphism $J : \mathcal{T}_g \longrightarrow \Lambda^3 H$:

$$J([f,h]) = J(fhf^{-1}) - J(h) = (f_* - 1)J(h) \tag{9}$$

Since J is onto and f_* can be any element of B_g, for $f \in \mathcal{L}_g^L$, this formula determines $J([\mathcal{L}_g^L, \mathcal{T}_g])$. We construct some examples.

1. Choose h so that $J(h) = x_i \wedge x_j \wedge y_k$ for i, j distinct, and f so that $f_*(y_k) = y_k + x_k$ and f_* is the identity on every other basis element. Then it is straightforward to compute, from Equation (9), that $J([f,h]) = x_i \wedge x_j \wedge x_k$.

2. Choose h so that $J(h) = y_i \wedge x_j \wedge y_k$ for i, k distinct, and f so that $f_*(y_i) = y_i + x_i$ and f_* is the identity on every other basis element. Then $J([f,h]) = x_i \wedge x_j \wedge y_k$.

3. Choose h so that $J(h) = y_i \wedge y_j \wedge y_k$, for i, j, k distinct, and f so that $f_*(y_i) = y_i + x_i$ and f_* is the identity on every other basis element. Then $J([f,h]) = x_i \wedge y_j \wedge y_k$.

4. Choose h so that $J(h) = y_i \wedge y_j \wedge y_k$, for i, j, k distinct, and f so that $f_*(y_i) = y_i + x_j$, $f_*(y_j) = y_j + x_i$ andf_* is the identity on every other basis element. Then

$$J([f,h]) = y_i \wedge x_i \wedge y_k + x_j \wedge y_j \wedge y_k + x_j \wedge x_i \wedge y_k$$

Now it is clear that the examples given here cover all the cases above.

This completes the proof of (b). In fact these arguments also prove (c) and so the proof of Theorem 1 is complete. □

3.4 Proof of Corollary 1

Recall the exact sequence (see e.g. [19])

$$1 \longrightarrow \pi_1(T) \xrightarrow{\eta} \Gamma_g \xrightarrow{e} \Gamma_{g,0} \longrightarrow 1$$

where T is the tangent circle bundle of $\Sigma_{g,0}$. The exact homotopy sequence of the bundle $T \longrightarrow \Sigma_{g,0}$ gives a central extension (since $\Sigma_{g,0}$ is orientable).

$$1 \longrightarrow \mathbb{Z} \longrightarrow \pi_1(T) \xrightarrow{\rho} \pi \longrightarrow 1$$

where $\pi = \pi_1(\Sigma_{g,0})$. η satisfies the following properties:

1. $\operatorname{Im} \eta \subseteq \mathcal{T}_g$

2. $\eta|\mathbb{Z}$ is defined by $\eta(1) =$ Dehn twist along a circle bounding the puncture, and $\eta|\pi$ is characterized by the property that the automorphism of π induced by $\eta(\alpha)$ is conjugation by α.

3. There is a commutative diagram

$$\begin{tikzcd}
\pi_1(T) \arrow[r, "\eta"] \arrow[d, "\rho"'] & \Gamma_g \arrow[d, "\xi"] \\
\pi \arrow[r, "\eta'"] & \operatorname{Aut}\pi
\end{tikzcd}$$

where $\operatorname{Aut}\pi$ is the group of automorphisms of π, $\eta'(g) =$ conjugation by g, and $\xi(f)$ is the automorphism of the fundamental group of $\Sigma_{g,0}$ induced by f.

Claim. (a) $\bar{J} \circ \eta = 0$

(b) *If* $\rho(\alpha) \in \pi$ *has homology class* $h \in H$*, then* $J_\omega \eta(\alpha) \equiv h \mod L$.

Assuming this, we then see, from (a), that $\bar{J}$ induces the homomorphism J_0 (since $\overline{\mathcal{L}^L_{g,0}} = e(\overline{\mathcal{L}^L_g})$) and, since, by (b), $J_\omega \circ \eta$ is onto (and $\mathcal{L}^L_{g,0} = e(\mathcal{L}^L_g)$), $\ker J_0 = \mathcal{L}^L_{g,0}$.

Proof of Claim. It follows from (3) above that $\hat{J}\eta(\alpha) \cdot x = h \wedge x$, for any $x \in L$, where $h \in H$ is the homology class of $\rho(\alpha)$. (In fact, since $\operatorname{Im}\eta \subseteq \mathcal{T}_{g,0}$, $\hat{J}(\eta(\alpha))$ is given by the standard Johnson homomorphism and this formula is true for any $x \in H$.) If $x \in L$, then $h \wedge x \longrightarrow 0 \in \Lambda^2(H/L)$– this proves (a). Similarly $J_\omega \eta(\alpha) \in \operatorname{Hom}(L, \mathbb{Z})$ is the functional $x \longrightarrow h \cdot x$ which, under the duality $\operatorname{Hom}(L, \mathbb{Z}) \cong H/L$, corresponds to the reduction of h. This proves (b). ☐

Finally note that $p(H) = 0$ since $p(h)$ is defined to be $\sum_i x_i \wedge y_i \wedge h$, for any symplectic basis $\{x_i, y_i\}$ of H and, since we can choose this basis so that $x_i \in L$, each term in this sum goes to $0 \in \Lambda^3(H/L)$. This completes the proof of Corollary 1. ☐

3.5 Proof of Corollary 2

Claim. *Suppose* $f' \in \Gamma_g$ *induces* $f \in \Gamma_{g,0}$*. Then* f'_* *(the induced automorphism of* F*) satisfies*

$$f'_*|F_L \equiv id \mod [F, F_L] \tag{10}$$

if and only if $f_*|\pi_L \equiv id \mod [\pi, \pi_L]$.

This follows from the fact that $\ker\{F \longrightarrow \pi\} \subseteq [F, F_L]$.

Now (10) is clearly equivalent to $f' \in \overline{\mathcal{L}^L_g}$ and $\bar{J}(f') = 0$, and this, by definition, is the same as $J_0(f) = 0$. So Corollary 2 is an immediate consequence of Corollary 1. ☐

4 Proof of Theorem 2

As pointed out above, we have already proved the last assertion of Theorem 2 and so we address the first part. Note that this is equivalent to the assertion that $J((\mathcal{L}^L_g)_m) = K_m$ for $m \geq 2$.

4.1 The case $m = 2$

We first show that $J((\mathcal{L}_g^L)_2) \subseteq K_2$. This is equivalent to the two inclusions:

1. $\Phi_f(H) \subseteq \ker\{\Lambda^2 H \longrightarrow \Lambda^2(H/L)\}$ and
2. $\Phi_f(L) \subseteq \Lambda^2 L$

where Φ_f is defined in Section 2.1. (1) is already proved in [7], so we address (2). According to Theorem 1, $\hat{J}$ induces a function $\mathcal{L}_g^L \longrightarrow \mathrm{Hom}(\hat{L}, X)$, where $X = \ker\{\Lambda^2 H \longrightarrow \Lambda^2(H/L)\}$. Now $\Lambda^2 L \subseteq X$ and the induced function $J' : \mathcal{L}_g^L \longrightarrow \mathrm{Hom}(\hat{L}, X/L^2 L)$ is a *homomorphism.* This follows from an alternative version of the crossed-homomorphism formula for $\hat{J}$:

$$\hat{J}(fg) = \hat{J}(f) + f_* \circ \hat{J}(g)$$

where $f_* : \Lambda^2 H \longrightarrow \Lambda^2 H$ is defined by $f_*(h_1 \wedge h_2) = f_*(h_1) \wedge f_*(h_2)$. If $f \in \overline{\mathcal{L}_g^L}$ then f_* induces the identity map on L and on H/L. From this it follows that $f_*|X \equiv 1 \mod \Lambda^2 L$ and so J' is a homomorphism. But then we can conclude that $J'((\mathcal{L}_g^L)_2) = 0$, which means that $\Phi_f(\hat{L}) \subseteq \Lambda^2 L$ for any $f \in (\mathcal{L}_g^L)_2$. This proves (2).

We now need to show that $K_2 \subseteq J((\mathcal{L}_g^L)_2)$. In fact we will show that $K_2 \subseteq J([\mathcal{L}_g^L \cap \mathcal{T}_g, \mathcal{L}_g^L]) = J([[\mathcal{L}_g^L, \mathcal{T}_g], \mathcal{L}_g^L])$. Let $\{x_i, y_i\}$ be an admissible basis of H. Then K_2 is generated by the following elements:

(a) $x_i \wedge x_j \wedge x_k$, for i, j, k distinct, and

(b) $x_i \wedge x_j \wedge y_k$, for $i \neq j$ and $j \neq k$.

Choose $f \in \mathcal{L}_g^L$ so that $f_*(y_i) = y_i + x_i$ and f_* is the identity on all other basis elements. By Section 3.3 we can choose $h \in \mathcal{L}_g^L \cap \mathcal{T}_g$ so that $J(h) = y_i \wedge x_j \wedge x_k$, if i, j, k are distinct. Then, by Equation (9), $J([f, h]) = x_i \wedge x_j \wedge x_k$. If $i \neq j$ and $j \neq k$, then, by Section 3.3, we can choose $h_2 \in \mathcal{L}_g^L \cap \mathcal{T}_g$ so that $J(h_2) = y_i \wedge x_j \wedge y_k$. Then by Equation (9), $J([f, h_2]) = x_i \wedge x_j \wedge y_k$.

This completes the proof of Theorem 2 for the case $m = 2$.

4.2 The cases $m \geq 3$

We first show that $J((\mathcal{L}_g^L)_m) \subseteq K_m$. We have already proved this for $m = 2$ in the previous section. We now argue by induction on m using Equation (9) and the fact that $(f_* - 1)K_m \subseteq K_{m+1}$. This latter inclusion follows from the expansion:

$$(f_* - 1)(a_1 \wedge a_2 \wedge a_3) = (f_* - 1)a_1 \wedge f_* a_2 \wedge f_* a_3 + a_1 \wedge (f_* - 1)a_2 \wedge f_* a_3 + a_1 \wedge a_2 \wedge (f_* - 1)a_3$$

and the following easy fact:

$$\text{If } f \in \overline{\mathcal{L}_g^L}, \text{ then } (f_* - 1)a \in L \text{ for any } a \in H, \text{ and } (f_* - 1)a = 0 \text{ if } a \in L \tag{11}$$

All that remains to prove is that $K_3 \subseteq J((\mathcal{L}_g^L)_3)$. But $K_3 = \Lambda^3 L$ is obviously generated by the elements $x_i \wedge x_j \wedge x_k$ and if we choose $h \in (\mathcal{L}_g^L)_2$ so that $J(h) = x_i \wedge x_j \wedge y_k$, which we can do by the previous section, and choose $f \in \mathcal{L}_g^L$ so that $f_*(y_k) = y_k + x_k$ and f_* is the identity on all other basis elements, then $J([f, h]) = x_i \wedge x_j \wedge x_k$

The proof of Theorem 2 is now complete. □

5 Proof of Theorems 3 and 4

The argument is a generalization of that in Section 4.2. Both these theorems will follow from:

Lemma 5.1. *1. If $f \in \Gamma_g$ and $g \in \Gamma_g[n]$, then $J_n([f,g]) = (f_* - 1)J_n(g)$.*

2. If $f \in \overline{\mathcal{L}_g^L}$, then $(f_* - 1)\mathcal{F}_n^r \subseteq \mathcal{F}_n^{r+1}$.

Proof of (1). This just follows from the easy formula $J_n(fgf^{-1}) = f_* J_n(g)$ and the fact that J_n is a homomorphism. □

Proof of (2). First we show that $(f_* - 1)\mathbb{L}_n^r(H) \subseteq \mathbb{L}_n^{r+1}$. $\mathbb{L}_n^r$ is generated by n-fold brackets $[a_1, \cdots, a_i \cdots, a_n]$, where at least r of the a_i belong to L . We can expand in the following manner:

$$(f_* - 1)[a_1, \cdots, a_i \cdots, a_n] = \sum_{i=1}^{n} [a_1, \cdots, a_{i-1}, (f_* - 1)a_i, f_* a_{i+1} \cdots, f_* a_n] \tag{12}$$

Now, it follows from 11 that each of the terms on the right side of 12 lies in $\mathbb{L}_n^{r+1}$.

Now suppose $a \otimes \alpha \in \mathcal{F}_n^r$. Then we can expand

$$(f_* - 1)(a \otimes \alpha) = (f_* - 1)a \otimes f_* \alpha + a \otimes (f_* - 1)\alpha \tag{13}$$

If $a \in L$ and $\alpha \in \mathbb{L}_n^{r-1}$, then the first term on the right side of 13 vanishes and the second lies in $L \otimes \mathbb{L}_n^r \subseteq \mathcal{F}_n^{r+1}$. If $\alpha \in \mathbb{L}_n^r$, then the first term lies in $L \otimes \mathbb{L}_n^r \subseteq \mathcal{F}_n^{r+1}$ and the second lies in $H \otimes \mathbb{L}_n^{r+1} \subseteq \mathcal{F}_n^{r+1}$. □

This completes the proof of the Lemma and, therefore, of Theorems 3 and 4.

6 Proof of Theorems 5, 6 and 7

6.1 Proof of Theorems 5 and 6

The framed pure braid group $\mathcal{P}_g$ on g strands is defined to be the group, under stacking, of pure braids, where each strand is equipped with a normal framing which is standard on the boundary. This is easily seen to be canonically isomorphic to $\mathcal{P}_g^0 \times \mathbb{Z}^g$, where $\mathcal{P}_g^0$ is the usual pure braid group, since there is a well-defined self-linking number for a normal framing. the imbedding $\psi : \mathcal{P}_g \longrightarrow \Gamma_g$ is defined in [24] as follows. Choose an imbedding of D_g, the disk with g holes, into Σ_g so that the internal boundary circles map to g independent meridians and the external circle maps to a bounding closed curve (see Figure 2).

According to the arguments of Artin, $\mathcal{P}_g$ can be regarded as the group of isotopy classes of diffeomorphisms of D_g which are the identity on the boundary. This representation of $\mathcal{P}_g$, together with the chosen imbedding $D_g \subseteq \Sigma_g$, defines ψ. Of course ψ depends on the choice of imbedding. An algebraic formulation of ψ can be given as follows. Let $\{x_i, y_i\}$ be a basis of $F = \pi_1(\Sigma_g)$ which satisfies:

1. The homology classes of $\{x_i, y_i\}$ form a symplectic basis of H
2. The boundary circle of Σ_g defines the element

$$(x_1 \cdots x_g)^{-1}(y_1 x_1 y_1^{-1} \cdots y_g x_g y_g^{-1}) \in F$$

See Figure 4. An element $\alpha \in \mathcal{P}_g$ is determined by a sequence $\lambda_1, \cdots, \lambda_g \in F'$, the free group on generators $x_1, \cdots, x_g$, which we shall call the *longitudes* of α. The longitudes satisfy the equation:



Figure 4. Definition of x_i and y_i

$$\lambda_1 x_1 \lambda_1^{-1} \cdots \lambda_g x_g \lambda_g^{-1} = x_1 \cdots x_g \tag{14}$$

Then $\psi(\alpha) \in \Gamma_g$ is the element which defines the following automorphism of F:

$$x_i \longrightarrow \lambda_i x_i \lambda_i^{-1}, \quad y_i \longrightarrow y_i \lambda_i^{-1} \tag{15}$$

Obviously ψ is an imbedding. It is not hard to prove the following Proposition (see [25]).

Proposition 6.1. *ψ induces an* imbedding *of the quotients:*

$$\psi_n : (\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} = \mathcal{P}_g[n]/\mathcal{P}_g[n+1] \longrightarrow \Gamma_g[n-1]/\Gamma_g[n]$$

for every $n \geq 1$.

$\{\mathcal{P}_g[n]\}$ is the weight filtration of $\mathcal{P}_g$, i.e. $\alpha \in \mathcal{P}_g[n]$ if and only if the longitudes $\lambda_i \in F'_n$. In fact $(\mathcal{P}_g)_n = \mathcal{P}_g[n]$.

Proof of Theorem 5. Since any diffeomorphism of D_g is a composition of Dehn twists along closed curves in D_g, then $\psi(\mathcal{P}_g) \subseteq \mathcal{L}_g^L$ if we set $L = \mathrm{Im}\{\mathrm{H}_1(D_g) \longrightarrow \mathrm{H}_1(\Sigma_g)\}$. Thus L is generated by $\{x_i\}$. Now suppose $\lambda_i \in F_n'$ are the longitudes of $\alpha \in (\mathcal{P}_g)_n$. Then $J_{n-1}\psi_n(\alpha) \in \mathrm{Hom}(H, \mathbb{L}_n(H))$ is given by $x_i \longrightarrow [\lambda_i, x_i] = 0 \in \mathbb{L}_n(H)$, $y_i \longrightarrow -\lambda_i$. The dual element of $H \otimes \mathbb{L}_n(H)$ is, therefore, $-\sum_i x_i \otimes \lambda_i$, which belongs to $L \otimes \mathbb{L}_n(L)$. ☐

Proof of Theorem 6. If $\alpha \in \mathcal{P}_g$ with longitudes $\lambda_i \in F'$, then we can write their reductions $l_i \in L = F'/F_2'$ in the form $l_i = \sum_j a_{ij}x_j$. It is clear that a_{ij} is the linking number of the i-th and j-th strands of (the closure of) α, if $i \neq j$, and the self-linking number of the i-th strand if $i = j$. Thus $A = (a_{ij})$ is a symmetric matrix and it is clear that any symmetric matrix can be realized by some pure braid. The definition of $\psi(\alpha)$ given in 15 shows that the symplectic automorphism determined by $\psi(\alpha)$ is represented by the matrix $\left(\begin{smallmatrix} I & -A \\ 0 & I \end{smallmatrix}\right)$. This proves (a).

To prove (b) we will construct an analogue of the Johnson homomorphism $J_b : (\mathcal{P}_g)_2 = \mathcal{P}_g[2] \longrightarrow \Lambda^3 L$ whose kernel is $(\mathcal{P}_g)_3 = \mathcal{P}_g[3]$ such that J_b induces an isomorphism $(\mathcal{P}_g)_2/(\mathcal{P}_g)_3 \otimes \mathbb{Q} \cong \Lambda^3 L \otimes \mathbb{Q}$. Suppose $\alpha \in (\mathcal{P}_g)_2$ has longitudes $\lambda_i \in F_2'$ and, under the isomorphism $F_2'/F_3' \cong \Lambda^2 L$, $\lambda_i \longrightarrow l_i$. We define $J_b(\alpha) = \sum_i x_i \otimes l_i \in L \otimes \Lambda^2 L$. It is clear that the kernel of J_b is $(\mathcal{P}_g)_3$. We see that $\mathrm{Im}\, J_b \subseteq \Lambda^3 L$ by a copy of Johnson's argument. Consider the exact sequence:

$$0 \longrightarrow \Lambda^3 L \longrightarrow L \otimes \Lambda^2 L \xrightarrow{\sigma} \mathbb{L}_3(L) \longrightarrow 0 \tag{16}$$

where $\sigma(a \otimes b) = [a, b]$. Equation (14), when read in F_3'/F_4', becomes $\prod_i [x_i, \lambda_i] = 1$ or, equivalently, $\sigma J_b(\alpha) = 0$.

Now the desired fact that $\mathrm{Im}\, J_b \otimes \mathbb{Q} = \Lambda^3 L \otimes \mathbb{Q}$ will follow from the fact that $d_g = \dim((\mathcal{P}_g)_2/(\mathcal{P}_g)_3 \otimes \mathbb{Q})$ is equal to $\dim(\Lambda^3 L \otimes \mathbb{Q})$. To prove this we use the split-exact sequence which computes the lower central series of the pure braid groups (see [2]):

$$1 \longrightarrow F_2'/F_3' \longrightarrow (\mathcal{P}_g)_2/(\mathcal{P}_g)_3 \longrightarrow (P_{g-1})_2/(P_{g-1})_3 \longrightarrow 1$$

From this sequence we get the recursive formula $d_g = d_{g-1} + \binom{g-1}{2}$. Solving this ($d_1 = 0$) gives $d_g = \binom{g}{3}$, the dimension of $\Lambda^3 L$. ☐

6.2 Proof of Theorem 7

First of all it is clear that any simple closed curve in D_g is a separating curve in Σ_g , under the imbedding $D_g \subseteq \Sigma_g$ which defines κ. Therefore the corresponding Dehn twist of Σ_g defines an element of $\mathcal{K}_g$. For the remainder of the proof we need an algebraic description of the map $\kappa : \mathcal{P}_g \longrightarrow \Gamma_g$. To do this we will find it convenient to introduce a map $\delta : F' \longrightarrow F$, where F' is the free group on generators$x_1, \cdots, x_g$ and F is the free group on $x_1, \cdots, x_g, y_1, \cdots, y_g$, defined by $\delta(x_i) = [x_i, y_i]$. Now suppose $\{\lambda_i\}$ are the longitudinal elements in F' which define the element $\alpha \in \mathcal{P}_g$. Then $\kappa(\alpha)$ is the element of Γ_g which corresponds to the automorphism of F given by

$$\begin{aligned} x_i &\longrightarrow \delta(\lambda_i)x_i\delta(\lambda_i)^{-1} \\ y_i &\longrightarrow \delta(\lambda_i)y_i\delta(\lambda_i)^{-1} \end{aligned} \tag{17}$$

If $\lambda_i \in F'_n$, then $\delta(\lambda_i) \in F_{2n}$ and so $\kappa(\alpha) \in \Gamma_g[2n]$. But it is proved in [24] that $\alpha \in (\mathcal{P}_g)_n$ if and only if $\lambda_i \in F'_n$ for every i. This shows that $\kappa((\mathcal{P}_g)_n) \subseteq \Gamma_g[2n]$.

For the proof of injectivity we will need the following:

Lemma 6.2. *δ induces an injection $F'_n/F'_{n+1} \longrightarrow F_{2n}/F_{2n+1}$.*

Proof. F'_n/F'_{n+1} is the free abelian group generated by the *standard basic Lie elements* $\{z_\nu\}$ on the symbols $x_1, \cdots, x_g$ (with that ordering) of degree n, as defined in [17, p. 334]. It will suffice, therefore, to observe that $\{\delta(z_\nu)\}$ are distinct standard basic Lie elements on the symbols $x_1, \cdots, x_g, y_1, \cdots, y_g$ (with that ordering) of degree $2n$. But it follows, by a straightforward inductive argument on the degree that these elements are basic and, in the *standard ordering*, as defined in [17, p. 334], $z_\nu < z_\mu$ implies $\delta(z_\nu) < \delta(z_\mu)$. □

To prove that κ_n is injective we use the Johnson-Morita map $J_{2n} : \Gamma_g[2n] \longrightarrow H \otimes \mathbb{L}_{2n+1}(H)$ whose kernel is $\Gamma_g[2n+1]$. If $\alpha \in (\mathcal{P}_g)_n$, then, from (17), we have

$$J_{2n}(\alpha) = \sum_i (x_i \otimes [\delta(\lambda_i), y_i] - y_i \otimes [\delta(\lambda_i), x_i])$$

It follows from this that if $J_{2n}(\alpha) = 0$, then $\delta(\lambda_i) \in F_{2n+1}$ for all i. From Lemma 6.2 we conclude that $\lambda_i \in F'_{n+1}$ and so $\alpha \in (\mathcal{P}_g)_{n+1}$.

Finally, to see that the induced map $(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \longrightarrow (\mathcal{K}_g)_n/(\mathcal{K}_g)_{n+1}$ is injective it suffices to observe that $(\mathcal{K}_g)_n \subseteq \Gamma_g[2n]$, for every n (see [23, Cor. 3.3]), and so the injection κ_n can be factored

$$(\mathcal{P}_g)_n/(\mathcal{P}_g)_{n+1} \longrightarrow (\mathcal{K}_g)_n/(\mathcal{K}_g)_{n+1} \longrightarrow \Gamma_g[2n]/\Gamma_g[2n+1]$$

□

7 Proof of Theorem 8 and Theorem 9

7.1 Proof of Theorem 8

Since $\mathcal{K}_g$ and $\mathcal{T}_g$ are normal and any two choices of L give conjugate subgroups $\mathcal{L}_g^L$ the truth of the assertions for any single choice of L implies the truth for any other choice. Recall from [8] the following material.

Let $i : \Sigma_{g,0} \longrightarrow S^3$ be any Heegard imbedding. and $L \subseteq H$ a suitable Lagrangian subspace. Then the association $f \longrightarrow S^3_f$, for any $f \in \overline{\mathcal{L}_g^L}$, defines a linear map $\mathbb{Q}\overline{\mathcal{L}_g^L} \longrightarrow \mathcal{M}$, where $\mathbb{Q}G$, for any group G, denotes the group algebra over $\mathbb{Q}$. This map induces the following three maps on the associated graded lower central series quotients:

$$\begin{aligned}
\phi_n^L &: \mathcal{G}_{3n}(\mathcal{L}_g^L) \otimes \mathbb{Q} \longrightarrow \mathcal{A}_n^{\mathrm{conn}}(\emptyset) \\
\phi_n^T &: \mathcal{G}_{2n}(\mathcal{T}_g) \otimes \mathbb{Q} \longrightarrow \mathcal{A}_n^{\mathrm{conn}}(\emptyset) \\
\phi_n^K &: \mathcal{G}_{n}(\mathcal{K}_g) \otimes \mathbb{Q} \longrightarrow \mathcal{A}_n^{\mathrm{conn}}(\emptyset)
\end{aligned}$$

where $\mathcal{A}_n^{\mathrm{conn}}(\emptyset)$ is the subspace of $\mathcal{A}_n(\emptyset)$ spanned by *connected* trivalent graphs and $\mathcal{G}_n(G)$ is the lower central series quotient G_n/G_{n+1}, for any group G.

It is proved in [8] that these maps are *onto* if $g \geq 5n+1$. Thus we can, for example, choose $f \in (\mathcal{K}_g)_n$ so that $\phi_n^K(f \otimes 1) \neq 0$. If $f \in (\mathcal{L}_g^L)_{3n+1}$ then we would have $\phi_n^L(f)$ defined and zero. But ϕ_n^L and ϕ_n^K take the same value on any element in both their domains, since they are defined by the same construction. Thus $f \notin (\mathcal{L}_g^L)_{3n+1}$. Similarly $f \notin (\mathcal{T}_g)_{2n+1}$.

The proofs of the other two non-inclusions are the same. □

7.2 Proof of Theorem 9

Let b, D and $K \subseteq S^3$ be as indicated on page 8. Let $N_0 \approx I \times D$ be a regular neighborhood of D in S^3 and N a regular neighborhood of $K \cup D$. Let $i : \Sigma_g \longrightarrow \partial N$ be a homeomorphism. Then $K_b \subseteq S^3$ is defined by replacing $K \cap N_0$, which consists of g straight line segments, by the pure braid b. Thus we have $K_b \subseteq N \subseteq S^3$. Let $D_g = D \times 1 \cap \Sigma_g$, which determines the homomorphism Ψ. See Figure 5.



Figure 5. The action of a pure braid on a knot

The main point here is the following:

Lemma 7.1. *There is a homeomorphism h of N onto itself which satisfies:*

1. $h(K) = K_b$,
2. $h|\partial N = i\psi(b)i^{-1}$.

Proof of Lemma. Recall the correspondence between a pure braid b and a homeomorphism h' of D_g onto itself. If b is considered as a path in the configuration space of g points in the 2-disk, then this path extends to a diffeotopy of the 2-disk. The end point of this diffeotopy fixes δD and the g points and h' is just the restriction to D_g. The isotopy class, rel δD of h' is uniquely determined. For *framed* pure braids, the same construction determines the isotopy class, rel δD_g, of h' h' is the identity on δD_g. The diffeotopy then defines a diffeomorphism $\bar{h}$ of $I \times D_g$ onto itself which is the identity on $0 \times D_g$, h' on $1 \times D_g$ and, if

d denotes the union of the g interior disks whose complement is D_g, $\bar{h}(I \times d) = b$. Here we think of a framed pure braid as a coordinatized tubular neighborhood of the actual braid. We can identify N_0 with $I \times D_g$ so that a tubular neighborhood of $N_0 \cap K$ is identified with $I \times d$. Then $\bar{h}$ becomes a diffeomorphism of N_0 and we can extend it over N by declaring it to be the identity on $N - N_0$. This defines the desired h. (1) is clear from the construction and (2) follows from the definition of ψ. $\square$

We now return to the proof of Theorem 9. Let N' be the closure of the complement of N in S^3. Then $S^3_{\psi(b)}$ is, by definition, $N \cup_f N'$, where $f = i\psi(b)i^{-1}$. But we can use the diffeomorphism h of Lemma 7.1 to construct a diffeomorphism $\tilde{h} : S^3_{\psi(b)} \longrightarrow S^3$ as follows. Regarding S^3 as $N \cup_{id} N'$ we define $\tilde{h}|N = h$ and $\tilde{h}|N'$ =identity. By Lemma 7.1, $\tilde{h}(K) = K_b$. The effect of $\tilde{h}$ on the framing of K is that an n-framing on K is mapped to a $(n + \sum_{ij} l_{ij})$-framing on K_b and so, for our choice of framing on b, a $+1$-surgery on $K \subseteq S^3_{\psi(b)}$ induces, via $\tilde{h}$, a $+1$-surgery on $K_b \subseteq S^3$.

Now $(S^3_{\psi(b)})_K = (S^3_K)_{\psi(b)}$ and when we show that $\psi(b) \in \Lambda_g^{L_j}$, for the given imbedding $j : \Sigma_g \subseteq S^3_K$, the proof of the Theorem will be complete. Clearly $\operatorname{Im}\psi \subseteq \Lambda_g^{L_i}$ where $i\Sigma_g \subseteq S^3$ is defined from K above. But $L_i \neq L_j$. In fact if $\{x_i, y_i\}$ is the symplectic basis defined by the meridians and longitudes of $\Sigma_g = \delta N-$ so that L_i is spanned by $\{x_i\}-$ then L_j is spanned by $\{x_i - y\}$, where $y = \sum_i y_i$. Thus to show $\psi(b) \in \Lambda_g^{L_j}$ we need to show that $\psi(b)(y) = y$. But if $\{l_{ij}\}$ is the linking matrix of the framed link b, then $\psi(b)(y_i) = y_i + \sum_j l_{ij}x_j$ and so $\psi(b)(y) = y + \sum_j(\sum_i l_{ij})x_j$. Since we have required $\sum_i l_{ij} = 0$ the proof is complete. $\square$

References

[1] D. Bar-Natan, *On the Vassiliev knot invariants*, Topology **34** (1995) 423–472.

[2] M. Falk, R. Randell, *The lower central series of a fiber-type arrangement*,

[3] S. Garoufalidis, *On finite type 3-manifold invariants I*, J. Knot Theory and its Ramifications **5**, no. 4 (1996), 441–462.

[4] S. Garoufalidis, N. Habegger, *The Alexander polynomial and finite type* 3*-manifold invariants.* preprint.

[5] S. Garoufalidis, J. Levine, *On finite type 3-manifold invariants II*, Math. Annalen, **306** (1996) 691–718.

[6] ———, *On finite type 3-manifold invariants IV: comparison of definitions*, Proc. Camb. Phil. Soc.

[7] ———, *Finite type 3-manifold invariants, the mapping class group and blinks*, Journal of Diff. Geom., to appear.

[8] ———, *Finite type 3-manifold invariants and the structure of the Torelli group I*, Inventiones Math., to appear.

[9] N. Habegger, *Finite-type 3-manifold invariants: a proof of a conjecture of Garoufalidis*, preprint.

[10] R. Hain, *Infinitesimal presentations of the Torelli groups*, Journal of the AMS, to appear.

[11] A. Hatcher, W. Thurston, *A presentation for the mapping class group of a closed orientable surface*, Topology **19** (1980) 221–237.

[12] D. Johnson, *An abelian quotient of the mapping class group*, Math. Ann. **249** (1980) 225–242.

[13] ______, *On the structure of the Torelli group II: A characterization of the group generated by twists on bounding curves*, Topology **24** (1085) 113–126.

[14] ______, *On the structure of the Torelli group III: the abelianization of $\mathcal{T}$*, Topology **24** (1085) 127–144.

[15] ______, *A survey of the Torelli group*, Contemporary Math. **20** (1983) 163–179.

[16] T.T.Q. Le, J. Murakami, T. Ohtsuki, *A universal quantum invariant of 3-manifolds*, Topology, to appear.

[17] W. Magnus, A. Karrass, D. Solitar, COMBINATORIAL GROUP THEORY: PRESENTATIONS OF GROUPS IN TERMS OF GENERATORS AND RELATIONS, Dover Publications, New York (1976).

[18] S. Matveev, M. Polyak, *A geometric presentation of the surface mapping class group and surgery*, Comm. Math. Physics **160** (1994) 537-556.

[19] S. Morita, *Characterisic classes of surface bundles*, Inventiones Math., **90** (1987) 551–577.

[20] ______, *Casson's invariant for homology 3-spheres and characteristic classes of vector bundles I*, Topology, **28** (1989) 305–323.

[21] ______, *On the structure and homology of the Torelli group*, Proc. Japan Academy, **65** (1989) 147-150.

[22] ______, *Mapping class groups of surfaces and three-dimensional manifolds*, Proceedings of ICM, Kyoto 1990, Springer-Verlag (1991) 665–674.

[23] ______, *On the structure of the Torelli group and the Casson invariant.* Topology **30** (1991) 603–621.

[24] T. Oda, *A lower bound for the graded modules associated with the relative weight filtration on the Teichmuller group*, preprint.

[25] T. Ohtsuki, *Finite type invariants of integral homology 3- spheres*, J. Knot Theory and its Ramifications **5** (1996) 101–115.

[26] T. Stanford, *Braid commutators and Vassiliev invariants*, Pacific Journal of Math.